MY PROTOCOLE FOR THE OIO

OBLIGATED INTRA-CELLULAR ORGANISMS




BILHARZIOSIS

IN ENGLISH

CÉCILE JADIN -JOHANNESBURG

COPYRIGHT RESERVED






AND AFTER ESTABLISHING THE DYSFUNCTIONS TO SINGLE OUT THE CARRIERS

FROM THE SICK ONES

- > FBC & ESR
- > Iron Study
- Kidney Functions
- Liver Functions
- Cholesterol
- > Glucose
- > T3 T4 Thyroids Antibodies
- > CRP, RF, ANF, CK, Cardiolipins
- Protein electrophoresis
- > IgE

TOGETHER WITH THE SYMPTOMS


BRIEF HISTORIC REMINDER ABOUT BORRELIOSIS




- o 7 DAYS /MONTH, as it takes the OIO 4 weeks to fill in an endothelial cell when they are active
- OF 2 ANTIBIOTICS, A BACTERIOSTATIC TO OPEN THE CELL,
- COMBINED WITH A BACTERICIDAL to kill what is inside the cell
- o ALTERNATED COMBINATION to avoid germs mutation,
- to avoid destroying normal cell, to allow normal intestinal bacteria to be restored
- o n.b. considering the modern alimentation antibiotics in meat and fish a daily intake of probiotics is recommended.
- The treatment also includes vitamin B complex as the antibiotics impair their normal production.
- A proton pump inhibitor to protect the stomach from the antibiotics is also prescribed.

IN THE PRESENCE OF BILHARZIOSIS







- Omnipresent on earth
- Needs a wee and a snail for transmission
- Laying from 200 to 2000 eggs/day
- Life span of 40 years
- Rarely causes hematuria
- More often epilepsy
- Kills more than 300.000 people/year
- Eggs will be retained in a pocket of calcium, mimicking a tumor, a cancer
- Gets through the blood brain barrier
- Needs at least 5 years of therapy

 praziquantel twice a year according to

OUR CHOICE OF ASSOCIATION OF ANTIBIOTICS FOR ADULTS:


- DOXYCYL 100MG X 2/DAY + CIPROFLOXACINE 500 MG X 2/DAY X7 DAYS- NO EXERCISE!
- LYMECYCLINE 300MG X 2/DAY X 7 + METRONIDAZOLE 400MG X 2/DAY X 7 DAYS NO ALCOHOL!
- MINOMYCINE 100MG X 2/DAY X 7 + RULIDE 150 MG X 2/DAY X 7 DAYS
- DOXYCYL 100 MG X 2/DAY + TAVANIC 250MG X 2/DAY X 7 DAYS- NO EXERCISE!
- CLACID 500MG X 2/DAY X 7 + AUGMENTIN 1.000 MG X 2/DAY X 7 DAYS
- DALACIN C150 MG 2 X2/ DAY X 7 DAYS GIVEN ALONE FOR BEST RESULTS
- DOXYCYL 100 MG X 2/DAY + AVELON 400MG X 2/DAY X 7 DAYS- NO EXERCISE!
- TETRALYSAL 400MG X 2/DAY + DAPSONE 100MG NOCTE (STOP IF BLUE LIPS) X7 DAYS
- VIBRAMYCINE 100MG X 3/DAY X 7 DAYS GIVEN ALONE FOR BEST RESULTS


ALL THOSE ANTIBIOTICS MUST BE TAKEN AFTER FOOD (NOT ONLY WATER)AND WITHOUT ANY DAIRY PRODUCTS.

PATIENTS MUST AVOID SUGAR INTAKE AND SUPPLEMENTS.

SPECIALLY: VITAMIN D AS THE COMPLICATIONS ARE MORE SEVERE THAN THE BENEFIT THROMBOGENICITY, CALCIUM DEPOTS ANYWHERE

Avoid the intake of Vitamin D3 - Calciferol





Our body requires Vit D1 from the sun

- O Avoid Statines. Cholesterol will normalise by itself during treatments
- O Avoid Iron supplements. This will feed the pathogens
- Avoid Cortisone, MTX, Biological treatments,
 Gammaglobulins. This will weaken any Immune System
- Support the thyroïd function (needed in 45 % of patients)
- N.B. the treatment will take from one to 3 years due to the nature of the OIO, going from active to dormant forms.
- Relapses are to be expected. About guarantied for Q Fever.